



Estimating the Lidar Ratios of Modeled Aerosol Species to Help Inform Selection for Marine Aerosol Lidar Ratios in a Future CALIPSO Data Products Release

Travis D. Toth¹ (travis.d.toth@nasa.gov), Gregory L. Schuster¹, Sharon D. Rodier², Marian Clayton², Mian Chin³, David Painemal², Zhujun Li², and Ellsworth Judd Welton³

¹NASA Langley Research Center, Hampton, VA

²Science Systems and Applications, Inc., Hampton, VA

³NASA Goddard Space Flight Center, Greenbelt, MD

103rd AMS Annual Meeting

12th Symposium on Lidar Atmospheric Applications

11 January 2023

The MAC Project for the MIRA Working Group



- **MIRA:** Models, In situ, and Remote Sensing of Aerosols
 - A new forum that encourages interdisciplinary work and fosters international collaborations
 - 200+ members in 22 countries and growing
 - Currently consists of 4 projects but are seeking new projects. If interested, please contact the MIRA steering committee at <https://science.larc.nasa.gov/mira-wg/contacts/>


- **MAC:** Mapping Aerosol lidar ratios for CALIPSO

- Lead: Greg Schuster (NASA LaRC)

MIRA webpage aerosol models to build global lidar ratio maps that can vary by season
These maps will be pertinent to the CALIPSO Version 5 extinction profile

MIRA news

Join us with these links!



<https://science.larc.nasa.gov/mira-wg/>


<https://espo.nasa.gov/lists/listinfo/mira>

Introduction & motivation for this study

- After 16+ years, the NASA CALIPSO mission will soon retire
- Narrow window of opportunity exists to improve quality of CALIPSO aerosol retrievals by advancing knowledge of aerosol lidar ratios for different aerosol types
- For CALIOP, the elastic backscatter lidar aboard CALIPSO, an aerosol lidar ratio is generally assumed to retrieve vertical profiles of aerosol extinction and subsequently column-integrated AOD


***Goal of this study:** creation of regional and seasonal climatological lidar ratio maps for marine aerosols by leveraging passive aerosol retrievals (MODIS) and global model simulations (GEOS/GOGART)

- Current CALIPSO lidar ratio selection process uses a single lidar ratio for each of the 7 CALIPSO tropospheric aerosol types, with large uncertainties for most aerosol types:




Data & methods

1. Collocate CALIOP backscatter profiles and MODIS AOD retrievals.
2. Perform Fernald (1972) inversion algorithm: use MODIS column-integrated AOD to constrain collocated CALIOP backscatter profiles and infer aerosol lidar ratios.
3. Use CALIPSO L3 Stratospheric Aerosol Profile Product to scale lidar ratios to account for stratosphere.
4. Create subset of data for only profiles with CALIOP-classified marine aerosols.
5. Compute sea salt volume fractions (< 2.5 km) from model data.
6. Collocate daily modeled sea salt volume fractions with Fernald-retrieved marine lidar ratios.
7. Develop empirical relationship between modeled sea salt volume fraction and Fernald-retrieved lidar ratios of marine aerosols.



MODIS AOD-constrained lidar ratios for CALIOP-classified marine aerosols (2006-2017)



Number	12,430,815
Minimum	0.000 sr
Maximum	117.95 sr
Mean	31.27 sr
Median	28.34 sr
Standard Deviation	16.24 sr

Sensitivity studies of retrieved lidar ratios

Aerosol Top Height

Aerosol Top Height	≤ 0.5 km	≤ 1.0 km	≤ 1.5 km	≤ 2.0 km
Number	616,498	5,511,503	10,003,988	12,065,528
Minimum	0.002 sr	0.001 sr	0.000 sr	0.000 sr
Maximum	117.53 sr	117.53 sr	117.53 sr	117.53 sr
Mean	37.00 sr	33.29 sr	31.54 sr	31.19 sr
Median	33.25 sr	29.85 sr	28.44 sr	28.25 sr
Standard Deviation	20.30 sr	17.84 sr	16.63 sr	16.22 sr


Horizontal Averaging

Horizontal Averaging	5 km	20 km	80 km
Number	1,214,672	4,389,575	1,998,067
Minimum	0.007 sr	0.002 sr	0.000 sr
Maximum	115.91 sr	117.09 sr	117.53 sr
Mean	23.05 sr	32.35 sr	38.71 sr
Median	21.57 sr	29.91 sr	35.07 sr
Standard Deviation	10.58 sr	16.02 sr	20.65 sr
Mean FMF	0.40	0.45	0.54
Mean SS VF	0.94	0.92	0.88

In this study, we consider aerosol top heights of ≤ 2.5 km and all horizontal averages

MODIS AOD-constrained lidar ratios & comparisons to past studies

12-year mean (2006-2017) on 1x1 degree grid




Study	Lidar Ratio (sr)	Method/Technique
Bohlmann (2018)	23	Raman
Dawson et al. (2015)	26	SODA AOD & CALIOP IAB
Masonis (2003)	25	<i>In situ</i>
Rittmeister (2017)	17	Raman
Rogers et al. (2014)	27	HSRL
Schmid (2003)	34	Fernald inversion
Voss (2001)	32; 36	Fernald inversion
Wang (2020)	30	Fernald inversion
Welton (2002)	33	Fernald inversion

Sea salt volume fractions from GEOS/GOCART

12-year mean (2006-2017) on 1x1 degree grid

Modeled Sea Salt Volume Fraction




Computation of Modeled Sea Salt Volume Fractions:

- Aerosol Species: SO_4 , SO_4v , NH_4a , NO_3 (3 bins), BC, BrC, OC, Dust (5 size bins), Sea salt (5 size bins), but excluding dust for this analysis
- Convert mass mixing ratios (kg kg^{-1}) to volumes (m^3)
- Below 2.5 km, sea salt volume fraction computed as:

$$\frac{\Sigma(\text{sea salt volume})}{\Sigma(\text{total aerosol volume, no dust})}$$


Using the model to help construct seasonal lidar ratio maps

Marine lidar ratios as a function of modeled sea salt volume fraction (2006-2017)



Seasonal analysis:
2006-2017 (JJA) lidar ratios (RSE $\leq 10\%$)

Constrained Lidar Ratio (532)



Summary & next steps

- Advantage of approach is using the model to fill gaps & help create climatologies that are consistent with MODIS AOD-constrained retrievals
- Maps will provide improved lidar ratio transitions in coastal regions
- Future efforts include:
 - Seasonal analyses
 - AeroCom multi-model mean for sea salt volume fraction
 - Two-layer aerosol lidar ratios (e.g., smoke over marine)
 - Other CALIOP tropospheric aerosol types
- Helps with final major data products release for CALIPSO (Version 5)
- Provides important aerosol lidar ratio information for future algorithm development of the next space-based lidars (e.g., the AOS elastic backscatter lidar: ALICAT on inclined orbit, AOS-I)

Acknowledgements

- Funding through support of the NASA CALIPSO Project
- CALIPSO data obtained from the NASA LaRC ASDC
- MODIS data obtained from NASA GSFC
- Model data obtained from NASA Discover server (NCCS)

ASDC


References





- Bohlmann, S., Baars, H., Radenz, M., Engelmann, R., & Macke, A. (2018). Ship-borne aerosol profiling with lidar over the Atlantic Ocean: from pure marine conditions to complex dust-smoke mixtures. *Atmospheric Chemistry and Physics*, 18(13), 9661-9679.
- Dawson, K. W., Meskhidze, N., Josset, D., & Gassó, S. (2015). Spaceborne observations of the lidar ratio of marine aerosols. *Atmospheric Chemistry and Physics*, 15(6), 3241-3255.
- Kim, M. H., Omar, A. H., Tackett, J. L., Vaughan, M. A., Winker, D. M., Trepte, C. R., ... & Magill, B. E. (2018). The CALIPSO version 4 automated aerosol classification and lidar ratio selection algorithm. *Atmospheric measurement techniques*, 11(11), 6107-6135.
- Li, Z., Painemal, D., Schuster, G., Clayton, M., Ferrare, R., Vaughan, M., ... & Trepte, C. (2022). Assessment of tropospheric CALIPSO Version 4.2 aerosol types over the ocean using independent CALIPSO-SODA lidar ratios. *Atmospheric Measurement Techniques*, 15(9), 2745-2766.
- Masonis, S. J., Anderson, T. L., Covert, D. S., Kapustin, V., Clarke, A. D., Howell, S., & Moore, K. (2003). A study of the extinction-to-backscatter ratio of marine aerosol during the Shoreline Environment Aerosol Study. *Journal of Atmospheric and Oceanic Technology*, 20(10), 1388-1402.
- Rittmeister, F. (2017). From the Caribbean to West Africa: Four weeks of continuous dust and marine aerosol profiling with shipborne polarization/Raman lidar—a contribution to SALTRACE.
- Rogers, R. R., Vaughan, M. A., Hostetler, C. A., Burton, S. P., Ferrare, R. A., Young, S. A., ... & Winker, D. M. (2014). Looking through the haze: evaluating the CALIPSO level 2 aerosol optical depth using airborne high spectral resolution lidar data. *Atmospheric Measurement Techniques*, 7(12), 4317-4340.
- Schmid, B., Hegg, D. A., Wang, J., Bates, D., Redemann, J., Russell, P. B., ... & Jefferson, A. (2003). Column closure studies of lower tropospheric aerosol and water vapor during ACE-Asia using airborne Sun photometer and airborne in situ and ship-based lidar measurements. *Journal of Geophysical Research: Atmospheres*, 108(D23).
- Voss, K. J., Welton, E. J., Quinn, P. K., Johnson, J., Thompson, A. M., & Gordon, H. R. (2001). Lidar measurements during Aerosols99. *Journal of Geophysical Research: Atmospheres*, 106(D18), 20821-20831.
- Wang, S. H., Lei, H. W., Pani, S. K., Huang, H. Y., Lin, N. H., Welton, E. J., ... & Wang, Y. C. (2020). Determination of Lidar Ratio for Major Aerosol Types over Western North Pacific Based on Long-Term MPLNET Data. *Remote Sensing*, 12(17), 2769.
- Welton, E. J., Voss, K. J., Quinn, P. K., Flatau, P. J., Markowicz, K., Campbell, J. R., ... & Johnson, J. E. (2002). Measurements of aerosol vertical profiles and optical properties during INDOEX 1999 using micropulse lidars. *Journal of Geophysical Research: Atmospheres*, 107(D19), INX2-18.

A cosmic background image featuring a dark space filled with numerous stars of varying brightness. Two prominent nebulae are visible: a blue one in the upper right and a green one in the lower right. A semi-transparent light blue horizontal band spans the middle of the image, serving as a backdrop for the title text.


Back up slides

Polyfit Lidar Ratio (532)






Lidar Ratio (S_a)



2006–2017 Mean Lidar Ratio (sr)




N



COUNTS



Relative Standard




Relative Standard Error (%)



$$\frac{\sigma/\sqrt{N}}{\langle S_a \rangle}$$


MODIS AOD



2006–2017 Mean Aqua MODIS AOD




Sea Salt Volume Fraction



2006–2017 Mean Sea Salt Volume Fraction



MODIS Fine Mode Fraction



2006–2017 Mean MODIS FMF



Using CALIPSO Level 3 Stratospheric Aerosol Product to scale the lidar ratios from constrained retrievals

- a. Find CALIOP/MODIS points that fall within each 5 x 20 degree latitude/longitude bin for each month and pull the stratospheric AOD
- b. Use the following equation to get the scaled lidar ratio that accounts for stratospheric aerosol loading:

$$S_{scaled} = \frac{\tau_{nostrat}}{\tau_{clmn}} S_{clmn}.$$

$$\tau_{nostrat} = \tau_{clmn} - \tau_{strat}$$

Title

- ?
- ?
- ?
- ?
- ?

Title

- ?
- ?
- ?
- ?
- ?